

Attorney docket number: 1822/117

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Application for Transfection of Human Embryonic Stem Cells

Invention of:

Nissim Benvenisty

Ofra Yanuka

Maya Schuldiner

Rachel Eiges-Avner

Attorneys:

Bromberg & Sunstein LLP

125 Summer Street

Boston, MA 02110-1618

Tel: (617) 443-9292

Fax: (617) 443-0004

Attorney Docket: 1822/117

TRANSFECTION OF HUMAN EMBRYONIC STEM CELLS.

Cross Reference to Related Applications

5 This application gains priority from provisional application 60/253,222, filed November 27, 2000 and 60/267,664 filed February 9, 2001, these applications being incorporated by reference herein.

Technical Field and Background Art

10 The present invention relates to preparations and methods of transfecting human embryonic stem cells, forming clonal preparations of pluripotent stem cells and enhancing a cell population in a human subject.

 It is known in the prior art that embryonic stem (ES) cells are pluripotent cells derived from the inner cell mass (ICM) of an *in vitro* fertilized embryo grown to the blastocyst stage. These cells are unique in their ability to grow indefinitely in culture while retaining a normal karyotype. Embryonic stem cells were first isolated from mice and were found to form aggregates or embryoid bodies *in vitro* which spontaneously differentiated into various cell types (Robertson, (1987) in Teratocarcinomas and Embryonic Stem cell, A Practical Approach, pp. 71-112). More recently, it has been shown that human embryonic stem cells can differentiate in culture to cells of the three germ layers that arise in initial stages of embryonic development (Itskovitz-Eldor, J. et al., (2000) Mol. Med., Vol. 6, pp. 88-95) and that their differentiation potential may be manipulated by the use of growth factors (Schuldiner, et al., (2000) Proc. Natl. Acad. Sci. U. S. A.). Various cell types may be identified in the differentiated human ES cells, e.g. neurons (Schuldiner, M., (2001) Brain res.), pancreatic β cells (Assady, S. et al., (2001) Diabetes, Vol.50, pp. 1691-1697), cardiomyocytes (Kehat, I. et al. J. Clin. Invest. 108, 407-414. (2001), and hematopoietic cells (Kaufman et al., (2001) Proc. Natl. Acad. Sci. U. S. A., Vol. 98, pp. 10716-10721).

 Embryonic stem cells (ES) provide a potentially unlimited supply of differentiated cells. The properties of stem cells that include self-renewal and vast differentiation potential are of major importance in such procedures as they serve as a solution for the overall lack of tissue sources. Studies in the mouse, rat and non-human primates have

shown that transplantations of embryonic and adult stem cells into tissues such as pancreas (Ramiya et al., (2000) Nat. Med, Vol. 6, pp. 278-282), brain (Isaacson et al., (1989) Exp. Brain Res., Vol. 75, pp. 213-220, Lee et al., (2000) Brain, Vol. 123, pp. 1365-1379, Deacon et al., (1998) Exp. Neurol., Vol. 149, pp. 28-41), spinal cord
 5 (McDonald et al., (1999) Nat. Med., Vol. 5, pp. 1410-1412), heart (Klug et al., (1996) J. Clin. Invest., Vol. 98, pp. 216-224) and bone marrow (Gutierrez-Ramos et al., (1992) PNAS, Vol. 92, pp. 9171-9175, Palacios et al., (1995) PNAS, Vol. 92, pp. 7530-7534) have to some extent resulted in penetration and normal differentiation in the host tissue followed, in some instances, by phenotypic improvement of the animal's health
 10 (McDonald et al., (1999).

Recently, human fetal brain cells were injected into the nigra-striatum of patients suffering from Parkinson's disease. The grafted cells caused improvement in most patients, though in some patients the transplantation caused serious adverse reaction probably due to uncontrollable proliferation of the fetal cells (Freed et al., (2001) N. Engl.
 15 J. Med., Vol. 344, pp. 710-719). Such deleterious results, alongside tumor formation are some of the clinical problems that may arise through the use of non-terminally differentiated cells that could possibly still proliferate in the host tissue. Safety steps are needed to deactivate or destroy cells that prove to be harmful to the patient.

In particular, human ES cells may be uniquely useful as an unlimited source of
 20 cells for the medical procedure of transplantation in numerous pathologies, and as a component in biomedical engineering as well as providing clues on early stages of human development. However, the clinical problems associated with transplantation and biomedical engineering should be addressed including problems associated with the reaction to the foreign cells by the host immune system, residual proliferation of the cells
 25 after transplantation and tumor formation or ectopic and excessive differentiation.

Genetic manipulation of differentiation of human ES cells would be desirable so as to obtain a uniform population of precursors or fully differentiated cells for in vivo or in vitro use.

30 **Summary of the Invention**

In a first embodiment of the invention there is provided a method of altering gene expression in a population of human embryonic stem cells that includes: introducing a polynucleotide into the population of cells, the polynucleotide containing a gene expression altering sequence so that gene expression in the embryonic stem cells prior to

introducing the polynucleotide is measurably different from gene expression after introducing the polynucleotide. In an example of the method the expression altering sequence is an enhancer for modulating gene expression in the population of embryonic stem cells. The expression altering sequence may be a gene encoding a protein, the protein not normally being expressed in the population of embryonic stem cells and the protein may be selected from a fluorescent protein and an antibiotic resistance protein. The fluorescent protein may be for example any of green fluorescent protein, lacZ, firefly Rennila protein, luciferase, red cyan protein and yellow cyan protein. The antibiotic resistance protein may be for example any of hygromycin, neomycin, zeocin and puromycin.

According to the above, introducing polynucleotides into cells may be facilitated by formulations that include a cationic lipid reagent, cationic non-lipid polymer transfection reagent, a liposomal transfection reagent for introducing into the population of cells. Alternatively, electroporation may be used.

In another embodiment of the invention, a method is provided for altering gene expression in a population of human embryonic stem cells; that includes: introducing into the population of cells by electroporation or in the presence of a cationic polymer, a DNA sequence corresponding to at least one of an enhancer, a promoter and a gene so as to alter gene expression in the population of embryonic cells in an amount to permit cells containing the DNA sequence to be distinguished from cells absent the DNA sequence.

According to this embodiment, the gene may encode a protein selected from a fluorescent protein, a suicide gene, a knockout protein and an antibiotic resistance protein. The promoter may be selected from rex-1, oct-4, oct-6, SSEA-3, SSEA-4, TRA1-60, TR1-81, GCTM-2, alkaline phosphatase, and Hes 1 promoters, the fluorescent protein may be selected from green fluorescent protein, lacZ, firefly Rennila protein, luciferase, red cyan protein and yellow cyan protein, the antibiotic resistance protein may be selected from hygromycin, neomycin, zeocin and puromycin, the suicide gene may be an inducible apoptic gene or encode a protein selected from herpes simplex thymidine kinase, inducible Diphtheria toxin, bacterial cytosine deaminase, and the knockout may causes the DNA sequence to be inserted into and render non-functional a gene encoding beta 2 microglobulin, HLA-1, HLA-2 or an INF receptor.

In another embodiment of the invention, a method is provided for purifying pluripotent embryonic stem cells from a heterogeneous population of cells, comprising: introducing into the cells, a DNA encoding a selectable marker under a promoter that is

specifically active in undifferentiated cell; separating those cells expressing the selectable marker from cells not expressing the marker; and obtaining purified pluripotent cells. Accordingly, the selectable marker may be a fluorescent marker for example, green fluorescent protein, lacZ, firefly Rennila protein, luciferase, red cyan protein or yellow cyan protein. The cells containing the marker may be separated from the cells lacking the marker using a fluorescent activated cell sorter or a laser scanning cytometer. Where the selectable marker is an antibiotic resistance marker, separating cells expressing the marker from those that cannot can be achieved by culturing the cells in a selective medium containing antibiotic.

10 In an embodiment of the invention, a method is provided for treating a human subject for a condition resulting from a deficiency of a selected cell type, that includes causing human embryonic stem cells to be transfected in vitro with a nucleic acid containing a marker under a tissue specific promoter; separating the selected cell type expressing the marker from cells not expressing the marker, and introducing the selected cell type into the subject so as to treat the condition. The nucleic acid may further contain a suicide gene. The suicide gene may be an inducible apoptic gene or encodes a protein selected from herpes simplex thymidine kinase, inducible Diphtheria toxin, bacterial cytosine deaminase. The marker may be a fluorescent marker or an antibiotic resistance protein. The fluorescent protein may be any of green fluorescent protein, lacZ, firefly Rennila protein, luciferase, red cyan protein and yellow cyan protein. The antibiotic resistance protein may be any of hygromycin, neomycin, zeocin and puromycin. The cells may be transfected by means of a cationic polymer transfection reagent including lipophilic or nonlipophilic reagents and liposomal reagents or by electroporation. In embodiments of the method, the nucleic acid may not contain viral genes. The selected cell type may be one of epidermal cells, dermal cells, muscle cells, cartilage cells, osteoblasts, osteoclasts, neurons, retinal cells, endodermal cells, hematopoietic cells, cells of the immune system or may be specialized cells from any functionally distinct organ in the human subject. The cell type may be administered to the subject by injection. Examples of conditions for treating with a selected cell type includes cancer, immune disorders, autoimmune diseases, diseases of aging, degenerative diseases including neurodegenerative diseases, and conditions associated with trauma.

In an embodiment of the invention, a cell population is provided that includes a substantially pure population of human embryonic stem cells containing an expression altering sequence of exogenous DNA.

In a further embodiment of the invention, a method is provided of producing a clonal pluripotent cell population from a mixture of pluripotent and differentiated cells; including: transfecting the mixture of cells in the presence of a cationic polymer or by electroporation with a DNA encoding a marker protein under a promoter that is

5 selectively active in cells of the inner cell mass of the embryo; and separating the embryonic stem cells from the differentiated cells according to the presence or absence of an expressed marker so as to produce the clonal pluripotent cell population. The promoter may be selected from rex-1, oct-4, oct-6, SSEA-3, SSEA-4, TRA1-60, TR1-81, GCTM-2, alkaline phosphatase, and Hes1 promoters. DNA transfection may occur in the

10 presence of a cationic polymer. The marker protein may be selected from a fluorescent protein and an antibiotic resistance protein. The fluorescent protein may be selected from green fluorescent protein, lacZ, firefly Rennila protein, luciferase, red cyan protein and yellow cyan protein, the antibiotic resistance protein may be selected from hygromycin, neomycin, zeocin and puromycin.

15 An embodiment of the invention includes a method for regulating cell viability of a population of cells in a subject, wherein the population of cells are derived from a human embryonic stem cell culture which has undergone directed differentiation, the method including introducing the population of cells into the subject, the population of cells containing an exogenous DNA encoding a suicide sequence, wherein the population

20 of cells are selected from the group consisting of: undifferentiated cells, partially differentiated or wholly differentiated cells; and treating the subject with an agent for activating the suicide sequence in the cells if the cells give rise to adverse events in the subject so as to cause cell death of the cells. For example, the dverse events may be hyperproliferation of the introduced cells.

25 In an embodiment of the invention, a method is provided for screening an agent to determine an effect on differentiation of cells *in vitro*, comprising: adding the agent to an *in vitro* cell culture of a population of genetically engineered human embryonic stem cells expressing a detectable marker under a cell specific promoter; providing the conditions for the embryonic stem cells to differentiate; and determining the effect on differentiation

30 of the agent. The detectable marker may be a fluorescent marker or an antibiotic resistant marker.

In an additional embodiment of the invention, a reagent cell population for supplying material for transplantation is provided consisting essentially of pluripotent human embryonic stem cells modified by foreign genetic material which is DNA not

normally present in embryonic stem cells; which occurs in embryonic stem cells but is not expressed in them at levels which are biologically significant; DNA which occurs in embryonic stem cells and has been modified so that it is only expressed by selected derivative cells; or any DNA that can be modified to be expressed by embryonic cells, derivative cells alone or in any combination thereof. For example, the foreign genetic material comprises genetic material encoding at least one selectable marker. At least one selectable marker may be a dominant selectable marker for example a gene encoding antibiotic resistance, a gene encoding a suicide protein, a fluorescent protein or an antibiotic resistant protein. The suicide gene may be hsv-tk and the suicide protein thymidine kinase, the fluorescent protein may be green fluorescent protein, lacZ, firefly Rennila protein, luciferase, red cyan protein or yellow cyan protein and the antibiotic resistance protein, hygromycin, neomycin, zeocin or puromycin.

Brief Description of the Drawings

The foregoing features of embodiments of the invention will be more readily understood by reference to the following detailed description, taken with reference to the accompanying drawings, in which:

Fig. 1 shows the efficiency of different techniques to introduce DNA into human embryonic stem cells. The scale bar is 100µm. Inset :human ES cells transiently transfected with EGFP under the control of the housekeeping gene E1F (elongation factor 1) promoter. Green fluorescent ES cellss incorporate the foreign DNA.

Fig. 2 shows transfection of human embryonic stem cells (a) transient transfection; and (b) stable transfection.

Fig. 3 shows electroporation of human embryonic stem cells.

Fig. 4 shows the expression of a transfected marker of undifferentiated cells as a method to identify and isolate pluripotent human embryonic stem cells. (a) Human ES cells underwent stable transfection with EGFP fused to murine Rex 1 minimal promoter sequence. The transfected ES cells and their differentiated cell derivatives are shown: simple embryoid body (sEB) and mature embryoid body (maEBs). The left and middle columns are photos of bright bright and dark fields respectively. The right column is the overlay of the two photos. Note that only the undifferentiated cells are fluorescent. The fluorescent ES colony is surrounded by differentiated non fluorescent cells. The simple EB is labeled only in the middle and nt in the peripheral primitive endodermal cells. Mature EBs are generally not fluorescent and only very distinct areas in them are

fluorescent (probably residual fluorescent cells). The scale bar indicates 100 μ m. (b) The stable transfection of human ES cells with constitutively expressed EGFP construct, driven by mouse PGK promoter. Overlay photos of the dark on bright field of the transfected ES cells (ES) and their differentiated cell derivatives are shown: simple embryoid bodies (sEB), mature embryoid bodies (mEB) and differentiating embryonic cells derived from dissociated embryoid bodies (DE). Note that GFP is expressed by all cells differentiated and undifferentiated in the proliferating ES colony as well as by all cells of simple and mature Ebs (including those in the outer layer of the sEB, where differentiation of primitive endoderm is taking place in the mouse EB. The scale bar is 100 μ m.

Fig. 5 (A) shows FACS analysis of Human ES cell transfected with Rex1-EGFP construct according to their intensity of green fluorescence emission. Cell samples of MEF (feeder layer) and undifferentiated human ES (grown in the presence of feeder cells) were used as controls. Fluorescent intensity was then compared between undifferentiated human ES cells, transfected human ES cells and their differentiated cell culture derivatives (obtained by growth in non-supplemented gelatin-coated plates). (A-1) MEF, (A-2) undifferentiated ES cells and MEF, (A-3) undifferentiated transfected ES cells plus MEF; and (A-4) partially differentiated transfected ES cells on gelatin.

Fig. 5 (B) shows cells sorting of three GFP-expressing cell lines performed by FACS. Following trypsin digestion, cell samples (3-4 replicates per clone) were evaluated for percentage of cell viability (84%) and sorted according to intensity of green fluorescent emission. The collected cell samples (25,000-50,000) were redetermined for cell viability (86%) and replated on MEF^{neo+} culture dishes (2,500-8,000 cells per dish). Following growth in vitro, cell culture dishes were inspected and recorded for total number of proliferating human ES cells (20% \pm 4% [n=11]).

Fig. 5 (C) shows photos of fluorescent labeled proliferating human ES colonies (top, bright field and bottom, dark field) obtained 4 days after cell sorting by FACS.

Fig. 6 shows the effects of Ganciclovir on human ES Cells expressing the herpes simplex thymidine kinase gene. (A) Human ES cell clones expressing the HSV-TK gene showing sensitivity to the presence of the pro-drug Ganciclovir (Ganc) (B) Dose response of HSV-TK+ clones to Ganciclovir treatment. Six clones (TK+) and human ES cells as controls were grown in presence of various concentrations of Ganciclovir. The bars represent Standard error values (C) Time course of the effects of Ganciclovir on HSV-TK+ cells. Six HSV-TK+ clones and control human ES cells were treated with 2x10⁻⁶M

Ganciclovir for 9 days. The experiment was repeated 4 times. After 9 days Ganciclovir was removed from the media and cells were grown for additional 10 days. The bars represent Standard error values.

Fig. 7 shows selective ablation of HSV-TK+ cells (A) Analysis of mixed cultures of GFP expressing cells (GFP+) with HSV-TK+ (TK+) cells. Following treatment with Ganciclovir selective elimination of the HSV-TK+ cells is shown by FACS analysis of GFP expression.

Fig. 8 shows sensitivity of teratoma cells to Ganciclovir (A) Differentiated HSV-TK+ teratoma cells showing sensitivity to the presence of the pro-drug Ganciclovir (Ganc.) (B) Time course of differentiated HSV-TK+ teratoma cells in response to 2×10^{-6} M Ganciclovir during 8 days. After 8 days Ganciclovir was removed from the media and cells were grown for additional 8 days. The bars represent Standard error values.

Fig. 9 Retention of normal karyotype and differentiation potential by the hsTK+ cell lines (A) Karyotype of an HSV-TK+ clone using G-banding. All cell lines have a normal, XX chromosome set (B) RT-PCR analysis of differentiation markers from the three germ layers on human ES cells, hsTK+ embryoid body (EB) and teratoma (Ter) cells. The markers used were: α -fetoprotein (α FP), albumin, amylase, β -globin, cardiac actin (cActin), enolase, keratin, glial fibrillary acidic protein (GFAP) and neurofilament heavy chain (NF-H). As control for presence of cDNA the housekeeping genes β -actin and glyceraldehyde-3-phosphate hydrogenase were used.

Detailed Description of Specific Embodiments

Definitions. As used in this description and the accompanying claims, the following terms shall have the meanings indicated, unless the context otherwise requires:

“Transfection” is the introduction of nucleic acid into cells in a population.

Transfection may occur *in vivo* as well as *in vitro*. One result of transfection is to produce a genetically engineered cell.

“Vector” is any commercial vector such is available in the market, for example, vectors from Clontech, Promega, In Vitrogen or New England Biolabs.

“Suicide sequence” in a cell is any DNA which when activated as a result of an externally administered agent acting either directly on the DNA, RNA or on protein expressed by the DNA results in apoptosis or damage to the cells containing the suicide sequence. Suicide genes can be under a constitutive promoter or a tissue specific promoter for example an ES cell specific promoter. When the cells are transplanted into

a subject *in vivo*, the externally administered agent may be provided orally or parenterally including by subcutaneous, intramuscular or intravenous injection or by transdermal means. Examples of suicide genes are inducible apoptotic genes and those encoding thymidine kinase, bacterial cytosine deaminase, inducible Diphtheria toxin.

5 “Transcription factors” refer to any of DNA, RNA or protein that become associated with the DNA to be transcribed and modulate the extent of transcription. Examples of transcription factors include hepatic nuclear factors (such as HNF3), muscle specific factors (such as MyoD), hematopoietic factors, pancreatic factors (such as PDX-1), homeobox genes and others known in the art.

10 “Cell specific promoters” are generally non-coding nucleotide sequences located upstream from a gene and regulate the tissue specific expression. For example, neuronal specific promoters are located upstream from encoding sequence for neurofilament heavy chain; cardiac promoters determine expression of cardiac proteins and actins in cardiac muscle cell, hematopoietic promoters determine expression of globin proteins including
15 beta globin, a liver promoter regulates expression of albumin in hepatocytes, and a pancreatic promoter regulates expression of insulin. Other examples of promoters are those that regulate expression of nestin, tyrosine hydroxylase, dopamine beta
hydroxylase, CD34, PGX-1, albumin, ISL-1 and ngn-3. These examples are not intended to be limiting.

20 “Markers” are DNA, RNA or protein that can be readily detected in cells and provide a means of distinguishing those cells containing the marker from those that lack the marker. Markers can be used to track cellular events in circumstances involving a changing environment. Markers can be intrinsic in the cells of interest or may be foreign and introduced into the cells to express proteins. For example, where foreign DNA
25 encodes markers these are sometimes called reporter genes. “Reporter genes” are those genes that “report” the presence of particular cells and may include cell specific enhancers and promoters that control whether tissue specific expression of a gene occurs and how it is modulated. Reporter genes may be introduced into cells by transfection. Transfection of cells with genes encoding reporter proteins provides a means for tracking
30 cells. Examples of reporter genes include green fluorescent protein, Lac Z, firefly Rennila protein, red, yellow or blue cyan fluorescent proteins or other fluorescent protein, including those found in marine animals. Other markers include antibiotic resistance proteins to protect cells against for example, neomycin, hygromycin, zeocine and puromycin.

“Expression altering sequence” includes any exogenous nucleic acid which when introduced into target cells either extra-chromosomally or chromosomally, has the ability to modulate gene expression in the same cell or by association, in other cells, by either enhancing or suppressing expression of proteins already being made in the cell or by causing a protein to be expressed that would not normally be expressed but which is naturally encoded by DNA in the cell, or by causing a protein to be expressed that would not normally be expressed in the cells and is not naturally encoded by DNA in the cell such as the protein from one species in cells from another species. Examples of expression altering sequences include: enhancers, promoters, transcription activators and genes as well as sequences for enhancing recombination including homologous recombination, between proximate DNA molecules. Cells expressing particular genes in suitable quantities may be used in cell therapy in a subject to correct defective gene expression associated with a condition in the subject. Examples of therapeutically beneficial proteins expressed by genes in differentiated cells derived from human embryonic stem cells include growth factors such as epidermal growth factor, basic fibroblast growth factor, glial derived neurotrophic growth factor, nerve growth factor, insulin-like growth factor (1 and 11), neurotrophin-3, neurotrophin -4/5, ciliary neurotrophic factor, AFT-1, lymphokines, cytokines, enzymes-for example, glucose storage enzymes such as glucocerebrosidase, tyrosine hydroxylase.

“Condition” describes a state that is manifest as different from normal and for which a human subject may seek treatment. Examples of conditions include cancers such as late stage cancers including ovarian cancer and leukemia, diseases that compromise the immune system such as AIDS and autoimmune diseases such as multiple sclerosis, diabetes mellitus, inflammatory bowel diseases such as Crohn’s disease, systemic lupus erythmetosus, psoriasis, rheumatoid arthritis, autoimmune thyroid disease and scleroderma, conditions affecting the nervous system such as muscular dystrophy, Alzheimer’s disease, Parkinson’s disease, spinal cord injuries, liver diseases such as hypercholesterolemia and other conditions for which replacement of damaged tissue is desirable such as in heart disease, cartilage replacement, burns, foot ulcers, gastrointestinal diseases, vascular diseases, kidney diseases, urinary tract disease and aging related diseases and conditions. The condition may be associated with defective genes, e.g. defective immune system genes, cystic fibrosis genes or other genetic diseases.

We have established herein methods and compositions for genetically engineering human ES cell lines and we describe an efficient protocol for transfecting these cells. By introducing genetic modifications into their genome, we can manipulate these cells *in vitro*, we can purify stem cell clones using selectable markers and we can use, track, manipulate and if desired, kill engineered stem cells in cell based therapies as well as for other biomedical and research purposes.

Markers can be used to isolate specific cell types from a heterogeneous culture. For example, when a population of cells is transfected with a DNA containing a gene that codes for a drug resistance and is driven by a tissue specific promoter then regardless of the cell types that contain the gene, the only cell which will survive in the presence of the drug is the cell type capable of expressing the drug resistance gene.

Human embryonic stem cells which have been genetically modified as described below may be used for many purposes not only related to learning more about the biology of differentiation but also for cell therapy in human subjects. Such uses include the following: (a) tracking the steps of differentiation of the embryonic stem cells, (b) identifying factors that affect differentiation and self renewal, (c) separating pluripotent stem cells from partially differentiated stem cells, (d) characterizing and selecting different types of differentiated cells, (e) isolating pure preparations of differentiated cells, (f) improving the yield of selected differentiated cells, (g) monitoring transplanted cells *in vivo* including monitoring cell movement, proliferation, and location *in vivo*, (h) destroying transplanted cells undergoing hyperproliferation *in vivo* including conversion to teratocarcinomas and (i) eliminating MHC related molecules to avoid transplant rejection.

Although ES cell lines are now available for a large array of mammalian species (including hamster, mink, pig, rabbit, sheep, cattle, common marmoset and rhesus monkey (for a review see Prell et al., (1999) Cell Tissues Organs, Vol. 165, pp. 220-236) and for humans (US Application Serial No. 09/918,702) these ES cell lines other than those of mice have not been transfected with exogenous DNA. In murine ES cells, electroporation was found to be the method of choice for introducing foreign DNA into ES cells (Thomas et al., (1987) Cell, Vol. 51, pp. 503-512). In contrast to mouse embryonic stem cells, we found that although human ES cells can be transfected by electroporation, improved results were obtained by transfection in the presence of cationic polymers.

We have tested a variety of chemical based methods for introducing exogenous DNA into human ES cells. These include cationic lipid reagents, non-liposomal formulations, and cationic polymer transfection reagents. Examples of these categories of reagent include lipofectamine (cation lipid), electroporation, FuGENE (non-liposomal formulations) and ExGen (cationic polymers including preparations of linear polymers of ethyleneimine).

Example 1 and Figure 1 shows the results of using Firefly Renilla Luciferase (luc) reporter gene, driven by the Herpes Simplex TK promoter (Dual Luc Reporter Assay Kit, Promega) introduced into growing colonies of human ES cells, either by LipofectAMINETM (cationic lipid) (Life Technologies, Bethesda, Md), FuGENETM 6 (non-liposomal formulation) (Boehringer Mannheim, Germany) or ExGenTM 500 (cationic polymer reagent) (Fermentas, Hanover, Md), according to the manufacturer's protocols. After 48 hours, transfection efficiencies were evaluated by measuring the relative activity of the enzyme in the presence of its substrate, in respect to protein concentration. A clear difference in transfection efficiency could be demonstrated between ExGen 500, as compared to FuGENE 6, LipofectAMINE and electroporation. Transfection with ExGen 500 delivered DNA into human ES cells with an increased efficiency of an order of magnitude over other reagents (Figure 1).

The introduction of markers into embryonic stem cells according to the specificity of the promoter permits the separation of transfected cells from non-transfected cells or the separation of one type of transfected cell from other transfected cells. Markers can include a novel physiological feature for example a feature that enables transfected cells to grow where they could not otherwise, an example being a drug resistance gene such as Neo which confers cytoprotection to cells growing in G418. Other types of markers may facilitate the physical separation of different cell types from a mixture using biochemical analysis (protein assays, enzymatic assays, receptor binding assays) or immunological assays. Where the marker is an optical or laser recognizable feature, transfected cells expressing marker may be recognized either *in vivo* or *in vitro* according to whether the marker is expressed or not. Single or multiple marker genes may be introduced into cells. DNA encoding the markers may be associated with other genes forming fusion proteins or DNA that may specifically direct differentiation or modulate the amounts of a particular target protein produced by the differentiated cells. The expression of markers in transfected cells is regulated by promoter and enhancer sequences which result in production of amounts of protein.

We have obtained pure clones of human ES cells that are genetically modified so that their undifferentiated phenotype can be followed, and selected for *in vitro*. For example, a reporter gene, green fluorescent protein (EGFP) (see US Patent 6,316,181 for a description of green fluorescent protein marker), under the control of an ES-specific promoter was introduced into the human ES cells. By tagging the undifferentiated cells with green fluorescence, we monitored the differentiation status of the cells in culture during growth and propagation as well as following spontaneous and induced differentiation. We used the well-characterized promoter sequence of the mouse Rex-1 gene. Rex-1 is a retinoic acid-regulated zinc finger protein, which is expressed in undifferentiated ES and EC cells and in the inner cell mass (ICM) of day 4.5 mouse blastocysts (Hosler et al., (1993) Mol. Cell. Biol., Vol. 13, pp. 2919-2928, Rogers et al., (1991) Development, Vol. 113, pp. 815-824). Other promoter sequences suitable for selected expression in embryonic stem cells include Oct-4, Oct-6, SSEA-3, SSEA-4, TRA 1-60, TRA 1-81, GCTM-2, alkaline phosphatase, Hes 1 and other homeobox genes. In Example 4, Rex-1 promoter was used as a representative ES specific promoter that is rapidly down regulated upon differentiation of the embryonic cells. By introducing Rex-1-regulated selectable gene markers into human ES cells, we expressed these genes in a pluripotent-dependent fashion and determined the differentiation status of these cells in culture.

In another embodiment of the invention, a Rex-1-EGFP expression vector, which includes an SV40-Neo selectable marker, was constructed and delivered (by ExGen) into proliferating undifferentiated human ES colonies. The following day, cells were trypsinized and replated on a feeder of MitC-treated MEFNeo+, allowing the clonal propagation of transfected cells by G418 selection. By 14 days in culture, Neo resistant fluorescent labeled colonies were isolated and propagated for several passages, to allow the establishment of individual cell lines, derivatives of single transfected human ES cells (Example 1).

Of the 10 established cell lines, 4 were examined for pluripotency by growth under different culture conditions. When grown on feeder cells in the presence of leukaemia inhibitory factor (LIF) (to support undifferentiated growth), high expression of EGFP is detected in the small and densely packed cells of the undifferentiated colony. The fluorescent lightening overlaps well with the discrete margins of the colony and is absent in the periphery, where spontaneous differentiation is taking place (Figure 4). By growing the cells as a monolayer in the absence of a feeder layer and LIF, we show that

the labeling is highly correlated with cell morphology and does not exist in the large and loosely attached cells that grow beyond and apart from the colony. Moreover, when induced to differentiate by growing as cell aggregates in suspension culture, fluorescence gradually declines. Initially, in the outer surface of 4 day simple EBs, where a layer of primitive endodermal cells is formed, and later, if maintained to form cystic EBs (20 days in suspension), practically ceases, apart from few cores of undifferentiated cells (Figure 4).

We distinguished between populations of undifferentiated and differentiated human ES using a fluorescence-activated cell sorter (FACS) (Figure 5) or by other fluorescent activated methods such as laser scanning cytometry (Kamentsky et al., (1997) Acta Cytologica, Vol. 41, pp. 123-143) by magnetic methods (Dolan et al., (1999) US 5,985,153) or by other methods known in the art. Cell samples of mouse embryonic fibroblasts (MEF), undifferentiated human ES cells and undifferentiated and differentiated transfected cell lines have been characterized according to their fluorescent emission. A clear difference in fluorescent intensity exists between the undifferentiated cultures of untransfected and transfected cell lines. In addition, when comparing EGFP transfected human ES to their differentiated derivatives, a shift in emission is observed (Figure 5).

In an embodiment of the invention, we established genetically engineered cell lines of human ES, which express EGFP, under the control of mouse Rex-1 promoter. The ES-specific expression of this marker gene has allowed us to identify human pluripotent embryonic stem cells during their growth and differentiation *in vitro*. By growing the cells under different culture conditions we show a direct association between cell morphology and pluripotency: high EGFP expression in the densely packed cells of the undifferentiated colony is immediately turned off in the large and loosely attached cells that initiate differentiation. In addition, we demonstrate how pluripotent stem cells are rapidly eliminated in a developmentally regulated pattern, during the formation of EBs. This begins in the outer surface of simple EBs, where a layer of endodermal cells is formed, and proceeds by their elimination from most parts of the cystic EBs. These observations are similar to those in mice, where differentiation has been proven to be the cause for pluripotent stem cell extinction (Mountfort et al., (1998) Reprod. Fertil Dev., Vol. 10, pp. 527-533).

The uniform expression of Rex1-regulated sequence by the cells in the growing colony illustrates that these cells have not lost the features of the ICM during their long

cultivation and are thus truly pluripotent. Moreover, since these genetically engineered cell lines have been established by the clonal expansion of single transfected cells, the possibility of variation in the developmental potential of the cells in the undifferentiated colony can be ruled out.

5 We have developed a stem cell selection approach in an attempt to facilitate the maintenance of human ES cells *in vitro*. Currently, the available methods applied for this purpose involve the identification and isolation of single colonies under a dissecting microscope; however, these procedures are time consuming and labor intensive. As an alternative we present a method for purifying undifferentiated cells, by cell sorting of
10 fluorescent labeled cells, from mixed populations of cells. Similar selection of undifferentiated clones may be achieved by introducing into the cells a gene that enables drug selection (such as neo resistance gene) under the regulation of a human embryonic stem cell promoter such as Rex-1 or Oct-4 and maintaining the cells in the presence of the selection drug (such as the G418 antibiotic). By generating pure populations of
15 undifferentiated cells, as described above, we can avoid the loss of human ES cultures due to their spontaneous differentiation *in vitro*.

Introducing a cell-specific selectable marker to the genome of undifferentiated human ES cells provides a model for isolating specific cell types for transplantation from heterogeneous cell cultures obtained by induced differentiation. In addition, the positive
20 or negative effect of selected biological agents on directing differentiation of the stem cells to a desired cell type can be determined so as to maximize numbers of the desired cells for transplantation purposes. The ability of various biological agents to increase, decrease or modify in some way the number and nature of differentiated cells derived from embryonic stem cells may facilitate the effective use of embryonic stem cells as
25 reagents for transplantation. In summary, the ability to transfect human embryonic stem cells with exogenous DNA containing marker genes or master genes under tissue specific promoters provides new opportunities to analyze the effect of an agent or culture environment on the outcome of differentiation of embryonic stem cells. Moreover, the assay may be used to establish the effect of an agent or culture condition to enhance or
30 diminish an amount of a selected differentiated target cell type.

The potential use of human stem cells in transplantation and other cell based therapies require attention to the clinical problems that may arise upon residual proliferation of the cells after transplantation, causing tumor formation or ectopic and excessive differentiation. Human embryonic stem cells are a powerful tool for

transplantation medicine. These cells have the ability to proliferate indefinitely in culture while remaining pluripotent. In addition, ES cells may be genetically manipulated so as to evade the host immune system. Thus they may serve as a cell source that is both unlimited and widely applicable (reviewed in Solter, (2000) Nat. Rev. Genet., Vol. 1, pp. 199-207; Pera, (2001) Curr. Opin. Genet. Dev., Vol. 11, pp. 595-599). One of the risks in the use of stem cells is their ability to massively proliferate. This may become a problem if residual amounts of stem cells remain present in the grafted sample and take on inappropriate fates (Freed, (2001) N. Engl. J. Med., Vol. 344, pp. 710-71).

The methods described herein may be used to eliminate human ES prior to or after transplantation by negative selection, so as to avoid the risk of tumor induction. In these circumstances, cells expressing a gene product from exogenous DNA are caused to die when selection is applied. In addition, clinical complications that may arise from stem cell transplantation can be avoided by transplanting cells containing genes expressing a product that may kill cells in the presence of an administered therapeutic agent.

Examples of "suicide" genes include a tetracycline inducible form of the diphtheria toxin (Maxwell, (1986) Cancer Res., Vol. 46, pp. 4660-4664) and the bacterial cytosine deaminase (Pandha, (1999) J. Clin. Oncol., Vol. 17, pp. 2180-2189). Any "suicide" gene known in the prior art may be used for the negative selection of embryonic stem cells or their products *in vivo* or *in vivo* in the manner described herein. "Suicide" genes controlled by an ES specific promoter will allow elimination of the stem cell population only.

In an embodiment of the invention, we have genetically engineered human ES cell lines to constitutively express the herpes simplex thymidine kinase (HSV-TK) gene. Expression of the HSV-TK protein confers sensitivity to the FDA-approved drug Ganciclovir, allowing specific ablation of HSV-TK+ cells at concentrations non-lethal to other cell types. As expression of this gene causes sensitivity to the FDA approved pro-drug Ganciclovir (i.v. ganciclovir) it allows specific targeting of injected cells, allowing non-intrusive removal of grafts in case of unwanted side effects.

Ganciclovir (GCV), currently marketed by Roche, is an antiviral agent that interferes with DNA synthesis. The drug is phosphorylated by viral thymidine kinases, and undergoes further phosphorylation by cellular kinases. In its triphosphorylated form, the drug inhibits DNA polymerase by acting as a terminator in the synthesis of DNA, and induces apoptosis in cells.

Our results show that indeed, at a large range of Ganciclovir concentrations (10^{-5} - 10^{-7}), which do not affect normal cells, HSV-TK+ ES and tumor cells are eliminated. Moreover, blood concentrations in patients receiving the recommended intra-venous dose of 5-50mg/Kg (Pescovitz, (1999) Transpl. Infect. Dis. 1 Suppl., Vol. 1, pp. 31-34) should be in the range of 2×10^{-4} to 2×10^{-2} M. Plasma measurements of Ganciclovir following a daily dose of 5mg/Kg show that the actual steady state rate is higher than 10^{-6} M (Klatzmann, (1998) Hum. Gene. Ther., Vol. 9, pp. 2585-2594). This concentration is in the range that effectively killed all the cells in the in-vitro experiments. As reversion rates are low it seems that in the event of malignant transformation, treatment with the pro-drug may reduce tumor mass by more than 6 orders of magnitude, allowing faster elimination of the cells. The potential of these cells to be used in transplantation medicine is enhanced by the fact that they have retained the normal properties of ES cells. This is demonstrated by the fact that no karyotype changes have occurred during the manipulation process. This is important in that it reduces the risk of malignant transformation upon injection into hosts. Perhaps more importantly, we have shown that genetically modifying the cells has not changed their capacity to differentiate into a wide variety of tissues. The expression of markers for neuronal, cardiac and pancreatic cells gives hope that they may be used in the treatment of diseases such as neurodegenerations, cardiomyopathies and diabetes mellitus. In order to ascertain the safety of selective elimination of human ES cells *in-vivo*, cellular transplantation experiments using animal models such as diabetic mice or spinal cord injured rats must be performed.

Previous experiments with “suicide” genes have focused on the treatment of malignancies by the use of gene therapy. Most gene therapy research has focused on introduction of the HSV-TK due to its reduced side effects (no leakage of toxin and resistance of body cells). Advanced stage clinical trials are ongoing for several malignancies such as melanomas and glioblastomas with some success (Klatzmann, 1998). One of the main problems in gene therapy is the delivery system, a problem that does not exist while working with human ES cells. These cells can be grown indefinitely in culture allowing the production of pre-made genetically modified clones at need. Our results point to the ability to make human ES cells a safer reagent in transplantation medicine without sacrificing their pluripotent nature.

All references described in this description are herein incorporated by reference.

Examples

Example 1: Protocol for Transfection of human embryonic stem cells.

DNA was introduced into human embryonic stem cells using electroporation, or transfection with Lipofectamine plus (Invitrogen Life Technologies, Gruningen, The Netherlands), with FuGENE (BoehringerMannheim Mannheim, Germany) and with
 5 ExGen (Fermentas, Hanover, MD). An example protocol is provided on page 25.

Cell Culture: Human ES cells were grown on a feeder layer of mouse embryonic fibroblasts (MEF) and then transferred to gelatin coated plates and cultured further to reduce the number of murine cells in the culture. Differentiation into embryoid bodies (EBs) was initiated by transfer to petri dishes, where the embryoid bodies remained in
 10 suspension. (Schuldiner 2000) differentiated embryonic (DE) cells were formed by dissociating the EBs after 5 days and culturing them as a monolayer.

More specifically, human ES cells were obtained as described in Thomson et al., (1998) Science Vol. 286, pp. 1145-1147. Cleavage stage human embryos produced by in vitro fertilization (IVF) were obtained after the requisite approval process. The embryos
 15 were cultured to the blastocyst stage and inner cell masses were isolated. ES cell lines were isolated from the embryos and a cell line was selected which had a normal XX karyotype after 6 months of culture and could be passaged continuously over several months without undergoing a period of replicative crisis. The cells had a uniform undifferentiated morphology when grown on Mitomycin-C treated mouse embryonic
 20 fibroblast feeder layer (obtained from day 13.5 embryos) in 80% KnockOut™ DMEM medium (Gibco-BRL), supplemented with 20% KnockOut™ SR - serum-free formulation (Gibco-BRL), 1mM glutamine (Gibco-BRL), 0.1mM βmercaptoethanol (Sigma), 1% non-essential amino acids stock (Gibco-BRL), Penicillin (50units/ml) and Streptomycin (50 µg/ml), in the presence of 103 units/ml of leukemia inhibitor factor
 25 (LIF) (Gibco-BRL), and 4ng/ml of basic fibroblast growth factor (bFGF) (Gibco-BRL) (Schuldiner et al., 2000).

The undifferentiated cell cultures were induced to differentiate in vitro into EBs by omitting LIF and bFGF from the growth media and allowing aggregation in petri dishes (Schuldiner et al., 2000). Alternatively, undifferentiated cells were allowed to
 30 undergo spontaneous differentiation by growing as a monolayer on 0.1% gelatin (Merck) coated plates, in the absence of LIF and bFGF.

Teratomas were formed by injecting the cells into nude mice (Robertson, (1987), tumors were removed, dissociated and the cells grown on 0.1% gelatin coated plates with human ES cells media. Ganciclovir (Sigma G2536) was administered one day after

plating at the stated concentrations and media was changed with fresh Ganciclovir every two days.

Plasmid Construction: A Rex-1-EGFP expression vector was constructed by the deletion of the CMV promoter sequence from pEGFP-N1 (Clontech), and introduction of the mouse Rex-1 promoter sequence (700bp) into a HindIII restriction site.

Transient transfection: To determine the efficiency of DNA transfection by each of the above methods, cells were transfected with a construct of firefly Rennila protein under the control of the TK promoter. The cells were harvested 48 hours following transfection and luminosity of the Rennila protein was monitored using a luminometer.

Results are provided in the histogram of Figure 1 showing the relative activity of the transfected gene (luminosity units per mg protein). Each experiment was repeated three times and standard error is given in each experiment. As can be seen from Figure 1, the most efficient method for transfection of hES occurred when Exgen reagent was used.

Transfection and Establishment of Transgenic Cell Lines: Fully expanded and undifferentiated human ES cells underwent stable transfection with Rex-1-EGFP plasmid DNA (Rex-1 is a gene specific to undifferentiated ES cells (the Rex-1 gene) ExGen-500 transfection system (Fermentas) was used. Transfection was carried out in the human ES cells in 6-well plates on feeder cells, and was performed as described by the manufacturer's protocol. Specifically, the cells were incubated at room temperature with the transfecting agent for 10 minutes (2 µg of plasmid DNA plus 1 µl of ExGen 500 in 1 ml media per well), centrifugation at 1100rpm for 5 minutes and incubation at 37°C in a moist chamber for 45 minutes. Residuals of transfecting agent were removed by washing twice with PBS. The following day, the cells were trypsinized and re-plated on 10cm² culture dishes containing inactivated MEFNeo+. Two days after transfection G418 (200ng/ml) was administered to the growth medium, allowing the selective propagation of transfected cells in culture. By day 14, Neo resistant fluorescent-labeled colonies were identified by a fluorescent microscope. Single transgenic colonies were picked off by a micropipette, dissociated into small clumps of cells and transferred into a 2cm² (24-well) culture dish, on a fresh feeder of MEFNeo+. The cells continuously proliferated in the presence of G418 and formed a large number of expanding undifferentiated colonies, which express EGFP consistently high by all of the cells within the colony. Overcrowded cultures were trypsinized and propagated in 10cm² culture dishes for several passages to allow the establishment of individual cell lines, derivatives of a single transfected human ES cells.

Shown in Figure 4 are the transfected ES cells and their differentiation as a simple embryoid body (sEB), or as cystic embryoid bodies (cEB). In Figure 4, the left and middle rows are photos of light and dark fields respectively. The right row is the overlay of the two photos. Note that only the undifferentiated cells are fluorescent. The fluorescent ES clone is surrounded by differentiated non-fluorescent cells. The simple EB is labeled only in the middle and not in the peripheral primitive endodermal cells. Cystic EBs are generally not fluorescent and only very distinct area in them are still fluorescent (probably residual undifferentiated cells.)

FACS Analysis FACS analysis of Rex1-EGFP expressing cells was performed on a FACSCalibur system (Becton-Dickinson, San Jose, CA), according to cell size, granularity and fluorescent intensity. Undifferentiated human ES cells were used to set the background level of fluorescence. Transfected cells of undifferentiated (grown on MEF in the presence of LIF) and partially-differentiated (obtained by growth on gelatin in the absence of LIF) cell cultures were analyzed for fluorescence intensity as compared to background control.

Example 2 Transfection with DNA for directed differentiation.

Transcription factors that regulate differentiation of specific cells (master genes) are transfected into human embryonic stem cells according to the method described in Example 1. The expression of the transfected master gene should allow further differentiation of human ES cells in a regulated manner with a predetermined outcome.

Example 3 Transfection of embryonic stem cells with DNA encoding a cell specific marker.

Embryonic stem cells were transfected with a DNA encoding a cell specific marker linked to marker genes according to Example 1. Cells expressing the marker can be monitored in culture, and selected for or sorted by fluorescent activated cell sorting (FACS) to provide a purified preparation of a particular type of cell. Such a system allows analysis of cells such as neuronal cells (when using a neuronal specific enhancer such as the enhancer for neurofilament light chain gene); heart muscle cells (when using a cardiomyocyte specific enhancer such as the enhancer for alpha cardiac actin gene); liver cells (when using a hepatocyte specific enhancer such as the enhancer for albumin gene); or pancreatic cell (when using a pancreatic islet specific enhancer such as the enhancer for the insulin gene).

Example 4 Transfection of a human embryonic stem cells with a fluorescent protein to identify undifferentiated cells.

Embryonic stem cells were transfected with a DNA containing an enhancer specific to undifferentiated cells (such as the enhancer for Rex-1 or Oct-4) linked to marker genes, according to Example 1. Cells expressing the marker can be followed in culture, selected for or sorted by fluorescent activated cell sorting (FACS) to provide a purified preparation of undifferentiated cells (see Figure 2 for a colony of cells containing a fluorescent protein under an enhancer specific to undifferentiated cells).

Example 5: Creation of genetically modified clones.

Cell culture - Human ES cells (Thomson, 1998) and EBs were cultured as described (Schuldiner, 2000). Teratomas were formed by injecting ES cells into nude mice (Robertson, 1987), tumors were removed, dissociated and the cells grown on 0.1% gelatin coated plates with human ES cells media. Ganciclovir (Sigma G2536) was administered one day after plating at the stated concentrations and media was changed with fresh Ganciclovir every two days. Cell densities were followed by staining with Methylene Blue (Sigma) dissolved in 0.1M Boric Acid (pH=8.5) after fixation with 2.5% Glutaraldehyde. Color was extracted by 0.1N HCl and emission was read at 650nm.

Transfection and establishment of transgenic cell lines Plasmids introduced to the human ES cells were a PGK-EGFP plasmid (example 1) or a PNT plasmid (Tybulewicz et al., (1991) Cell, Vol. 65, pp. 1153-1163) that contains two PGK promoters driving either neomycin resistance gene or the herpes simplex thymidine kinase gene.

Transfection was performed using ExGen 500 (Fermentas) as described in Example 1.

FACS analysis and cell sorting - FACS analysis of PGK-EGFP and PNT expressing cells was performed on a FACSCalibur system (Becton-Dickinson, San Jose, CA), according to their green fluorescent emission.

RNA and RT-PCR -Total RNA was extracted as described (Chomczynski, P. and Sacchi, N. (1987) Anal. Biochem. 162, 156-159) and cDNA was synthesized from 1µg total RNA, using random hexamer (pd(N)6) as primer (Pharmacia Biotech) and M-MLV reverse transcriptase (Gibco-BRL). cDNA samples were subjected to PCR amplification with previously described DNA primers (Schuldiner, 2000) and primers for GFAP: SEQ ID. NO. 1: cgagaacaacctggctgcctatag and SEQ. ID NO. 2: gtgggtcctgcctcacatcacatc.

Cytogenetic analysis: Cells prepared for cytogenetic analysis were incubated in growth media with 0.1µg/ml of Colcemid for 10min, trypsinized, re-suspended in 0.075M

KCL and incubated for 10 min at room temperature, then fixed in 3:1 methanol/acetic acid and G-banded.

In order to create lines of cells carrying a negative selection marker we transfected human ES cells with a plasmid encoding the thymidine kinase gene from herpes simplex (HSV-TK) under control of a house-keeping gene (phosphoglycerate kinase-PGK) promoter. The introduced plasmid vector also contains the neomycin resistance gene that allowed selection for cells harboring the foreign DNA. Expression of the HSV-TK causes conversion of the pro-drug nucleoside Ganciclovir to its drug form as a phosphorylated base analog. The phosphorylated Ganciclovir is incorporated into the DNA of replicating cells causing irreversible arrest at the G2/M checkpoint followed by apoptosis (Halloran, (1998) Cancer Res., Vol. 58, pp. 3855-3865; Rubsam, (1998) Cancer Res., Vol. 58, pp. 3873-3882). Due to the high specificity of the cellular TK, Ganciclovir is not converted to the active form in normal eukaryotic cells causing insensitivity to its possible deleterious effects. All isolated lines of human ES cells transfected with HSV-TK were terminally arrested in response to the administration of Ganciclovir (Figure 6a). The transfected clones died in the presence of a range of Ganciclovir concentrations varying over three orders of magnitude (from $2 \times 10^{-8} \text{M}$ to $1 \times 10^{-5} \text{M}$), while the control human ES cells remained unaffected (Figure 6b). We chose to continue working at an intermediate concentration of $2 \times 10^{-6} \text{M}$ Ganciclovir, in which all HSV-TK+ cells died. At this concentration removal of Ganciclovir from the media after nine days of selection did not result in renewed growth of the cells, indicating that the action of the drug was terminal (Figure 6c).

Susceptibility of HSV-TK+ cells to Ganciclovir: Transplantation of human ES cells will expose them to conditions differing than those encountered in culture. First, they will differentiate over the passage of time and will also be surrounded by cells that are not affected by the treatment. Moreover, *in vivo* they will be free from the positive selective pressure of the neomycine to retain the plasmid and may be under strong negative selective pressure in favor of those cells that have lost the HSV-TK gene if treatment with Ganciclovir commences. These problems may cause resistance to the treatment suggested. In order to examine whether HSV-TK+ cells can be selectively eliminated, we have grown these cells together with cells expressing the green fluorescent protein (GFP+) and treated them with Ganciclovir. When the cultures were analyzed using a fluorescence activated cell sorter (FACS), it was clear that selective killing of the HSV-TK+ cells occurred at the same time scale and dose as with cultures containing

HSV-TK+ cells alone (Figure 7). Analysis of the effects of differentiation was performed by taking differentiated teratoma cells that have formed following injection of the clones into immunodeficient mice. The teratomas were removed from the mice after 8 weeks in which they were not subject to selection and the cells that were grown in culture retained their susceptibility to Ganciclovir at the same concentrations previously used (Figure 8a, b). This analysis also confirms that the fetal cells that are not grown continuously on neomycin selection still retain HSV-TK expression. In addition, we monitored the reversion rate of the Ganciclovir susceptibility in order to get an estimation of the possible number of drug resistant cells that arise following injection. The cells were grown for two passages in the absence of the positive selection drug neomycin, and then re-plated in the presence of Ganciclovir and grown for ten additional days and resistant colonies counted. Reversion was noted at a frequency between 10^{-6} - 10^{-7} .

Retention of normal karyotype and differentiation potential of modified clones:

Our aim was to ascertain that the genetically modified clones still retained their normal, pluripotent nature. Cytogenetic analysis of the transfected clones revealed a normal karyotype (Figure 9a). In addition, the cells could easily aggregate to form normal looking embryoid bodies (EBs) (Itskovitz-Eldor, 2000) *in vitro* and teratomas *in vivo*. RT-PCR analysis of cDNA from both EBs and Teratomas cells revealed a capacity to differentiate to cells from all three germ layers and to a wide variety of cell types as could be seen by the expression of the endodermal tissue specific markers: α -fetoprotein (primitive endoderm) albumin (liver) and amylase (pancreas); the mesodermal markers: β globin (blood), c-actin (cardiac muscle) and enolase (muscle); and the ectodermal markers: keratin1 (skin), GFAP-glial fibrillary acidic protein (Glia) and neurofilament (mature neurons) (Figure 9b).


Example 6 Immunological Tolerance in Transplanted Cells.

We target the genes of the major histocompatibility complex in embryonic stem cells and their differentiated progeny. This is achieved by knock-out or inhibition (by anti-sense or dominant negative form overexpression or ribozymes) of beta2 microglobulin or HLA-1 or HLA-11 or INF receptors. Any known method for inserting, deleting or modifying a desired gene from a mammalian cell combined with the transfection techniques described in Examples 1-5 can be employed. Methods and vectors for effecting gene knockout are the subject of numerous patents including US Patent 6, 074,853, 5,998,144, 5,948,653, 5,945,339, 5,925,544, 5,869,718, 5,830,698,

5,780,296, 5,614,396, 5,612,205, 5,468,629, 5,093,257 all of which are incorporated by reference in their entirety herein.

TOGETHER

Transfection of human embryonic stem cells



SEQUENCE LISTING

<110> Benvenisty, et al.

<120> Transfection of Human Embryonic Stem
Cells

<130> 1822/117

<160> 2

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 24

<212> DNA

<213> GFAP primer

<400> 1

cgagaacaac ctggctgcct atag

24

<210> 2

<211> 24

<212> DNA

<213> GFAP primer

<400> 2

gtgggtcctg cctcaatca catc

24

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
20